

**Home Care Instructions
Mitoxantone Treatment
(Patients Name and Date Here)**

<u>Clinical Signs</u>	<u>Monitor</u>	<u>Call Hospital () -</u>
Appetite	Picky, but still eats	Inappetence
Attitude	Slightly Lethargic	Lethargy and/or Reluctant to Move
Bowel Movement	Soft stool	Diarrhea
Temperature	<103 F	>103 F
Urination	Normal	Bloody Urine
Vomiting	Single Event	Frequent and/or Retching
Mucous Membrane	Pink/Moist	Pale/Jaundice
Eyes	Clear	Signs of Conjunctivitis
Right/Left Front Leg	Injection Site	Red/Swollen/Painful
Allergic Reaction	>>>>>>>>>>	Facial Twitching
Allergic Reaction	>>>>>>>>>>	Hives
Allergic Reaction	>>>>>>>>>>	Swelling Primarily on the Face

- Wear double layered gloves when cleaning up urine, feces, or vomit for 48 hours after receiving Mitoxantrone. Discard material in garbage bag and clean area thoroughly with regular cleaner and try to cover or avoid carpeted areas for 48 hours after cleaning. For pet waste outside, try to avoid the area for 48 hours and spray with water to dilute waste.
- A blue-green color to urine or a bluish color to the sclera may be noted but is of no concern.
- Maintain a low-stress environment during treatment.
- Ensure proper scheduling for follow-up laboratory work and next Mitoxantrone treatment.
- If at any time you have any questions or concerns please do not hesitate to call our office. No question is too small.

Owners Signature _____ Date _____

Released By _____ Date _____